



PRESS RELEASE

Monday 7 December 2009

Controversial kidney transplant technique could provide lifeline for very ill patients

Surgeons who successfully performed kidney transplants after removing small cancerous and benign masses from the donated organs, have published their results in the December issue of the urology journal **BJUI**.

The technique, carried out by US surgeons at the University of Maryland School of Medicine in Baltimore, could offer a vital lifeline to patients with end-stage renal disease as well as increasing the supply of viable organs.

“Transplanting a living donor kidney which has been affected by a renal mass is controversial and considered a high risk” says co-author Dr Michael W Phelan. “However the ongoing shortage of organs from deceased donors, and the high risk of dying while waiting for a transplant, prompted five donors and recipients to push ahead with surgery after the small masses were found in the donor kidneys.”

The five renal masses, which were discovered during routine donor evaluation, ranged from 1.0cm to 2.3cm in size. Cancerous cells were found in three of the five masses and the other two were benign.

The kidneys were removed from the donors, put on ice and taken to the recipients' operating rooms. Surgeons carefully removed the renal mass and a portion of the tissue near the mass was rushed through to pathology for confirmation that the tumour had been completely removed. The kidneys were then reconstructed and transplanted into the recipients.

One patient developed acute humoral rejection after surgery and was treated appropriately. There were no long-term problems in the transplanted kidneys and four of the patients were alive at the last follow-up, which ranged from nine to 41 months. The fifth died from an unrelated accident about a year after the transplant. None of the donors or recipients showed any evidence of recurring tumours.

The patients in the study ranged from 47 to 61 years of age, with an average age of 54, and the donors ranged from 38 to 72, with an average age of 38. Two of the five donor pairs were unrelated and three were genetically related. All the patients had end-stage renal disease, together with significant other illnesses, such as severe high blood pressure and complex heart problems.

Prior to the transplant, detailed discussions were carried out with each recipient and donor following the discovery of the renal masses in the donor kidney, so that they were both aware of the risks, including recurrence of the tumour.

“The global increase in patients with end-stage renal disease highlights the importance of identifying novel means to increase the donor pool” says Dr Phelan.

“Although donor transplants using organs from deceased people have risen 16 per cent and living donor transplants have risen by 68 per cent, there continues to be a significant shortage and many patients die each year while waiting for a transplant.

“The current study provides evidence to suggest that kidneys from donors with renal masses offer a minor, yet feasible, solution to the current organ shortage. These organs can be transplanted into recipients with limited life-expectancy on haemodialysis after careful removal of the renal mass. However, diligent follow-up of the donor and recipient is imperative in these cases.”

-ends-

For further information or a copy of the full paper please contact
Annette Whibley, Wizard Communications
0121 705 3575 / 07941 465757 (UK)
wizard.media@virgin.net

Notes to editors

- Living-donor renal transplantation of grafts with incident renal masses after ex-vivo partial nephrectomy. Sener et al. **BJUI**. 104, 1655-1660. (December 2009).
DOI: 10.1111/j.1464-410X.2009.08681.x
- Established in 1929, **BJUI** is published 23 times a year by Wiley-Blackwell and edited by Professor John Fitzpatrick from Mater Misericordiae University Hospital and University College Dublin, Ireland. It provides its international readership with invaluable practical information on all aspects of urology, including original and investigative articles and illustrated surgery. www.bjui.org
- **Wiley-Blackwell** is the international scientific, technical, medical, and scholarly publishing business of John Wiley & Sons, with strengths in every major academic and professional field and partnerships with many of the world's leading societies. Wiley-Blackwell publishes nearly 1,500 peer-reviewed journals and 1,500+ new books annually in print and online, as well as databases, major reference works and laboratory protocols.

For more information, please visit www.wileyblackwell.com or
www.interscience.wiley.com